

<i>Rubus occidentalis</i> L.		Black Raspberry.
Thickets and fence rows; occasional.		Thimbleberry.
Prunus emarginata Steyermark	May	
* <i>Rubus argutus</i> Link		Tall Blackberry
Sligo 1895.	May	
<i>Rubus cuneifolius</i> Pursh		Sand Black Berry
LEGUMINOSAE Sandy soil; occasional below the fall line.	June	
* <i>Rubus hispida</i> L.		Swamp Blackberry
Low woods; wet places.	June	
Witskie 1938. North Takoma 1912; Takoma Park 1895.		
* <i>Rubus procumbens</i> Muhl.	Dewberry	
<i>Rubus canadensis</i> of Ward's Flora	May	
Takoma 1938 Witskie; Terra Cotta, date not given.		
<i>Agrimonia</i> (Tourn.) L.		Agrimony
* <i>Agrimonia gryposepala</i> Wallr.		
Dry woods; rare.	August	
Takoma 1902.		
<i>Agrimonia mollis</i> (T. and G.) Britton		
<i>Agrimonia pubescens</i> Wallr.		
Open dry woods; occasional.	July	
<i>Agrimonia parviflora</i> Ait.		
Sandy alluvial soil; occasional	July-August	
<i>Rosa</i> (Tourn.) L.		Rose
<i>Rosa rubiginosa</i> L.		Sweetbrier. Eglantine
Rocky and open places; frequent.	June	
European.		
<i>Rosa carolina</i> L.		
<i>Rosa palustris</i> Marsh.		
Swamp borders and streams;	June	
occasional.		
<i>Rosa virginiana</i> Mill.		
Borders of swamps; wet places;	June	
occasional.		
<i>Prunus</i> (Tourn.) L.		Plum, Cherry, etc.
<i>Prunus serotina</i> Ehrh.		Wild Black Cherry
<i>Padus virginiana</i>		
Occasional.	May	

Prunus americana Marsh. Wild Plum
Occasional in thickets etc. April

Prunus persica (L.) Stokes Peach

Amygdalus persica L. Native of Asia.

LEGUMINOSAE (PULSE FAMILY)

Albizia Durazzini

Albizia Julibrissin Durazzini (Often called Mimosa)
Frequently cultivated as highly ornamental
tree. Rarely escaped.

Gleditsia Gleditsia L. Honey Locust

Gleditsia triacanthos L. Honey Locust
Frequent. May

Cassia (Tourn.) L. Senna

Cassia marilandica L. Wild Senna
Thickets etc.; common July-August

Cassia Chamaecrista L. Partridge Pea
Chamaecrista fasciculata (Michx.) Greene
Dry soil; common July-October

Cassia nictitans L. Wild Sensitive Plant
Chamaecrista nictitans (L.) Moench.
Sandy fields; common. July-October

Ceris (L.) Geris L. Redbud. Judas Tree

Ceris canadensis L. Rich woods; frequent. April

Baptisia Baptisia Vent False Indigo

Baptisia tinctoria (L.) R. Br. Wild Indigo
Dry woods; frequent.

Crotalaria (Dill) L. Rattle-box

Crotalaria sagittalis L. July-August
Sandy soil; not common.

Genista L. Wood-waxen. Whin.

**Genista tinctorici* L. Dyer's Greenweed
Rare. Fourteenth and Decatur 1913.
In Rock Creek valley.
Native of Europe and Asia.

<i>Lupinus</i> (Tourn.) D C.	Lupino
<i>Meibomia</i> Grisebach (Walt.) Kuntze	
* <i>Lupinus perennis</i> L.	Wild Lupine
Sandy soil; infrequent.	April-May
Takoma 1896.	
<i>Trifolium</i> (Tourn.) Sock	Clover. Trefoil.
<i>Meibomia</i> <i>Trifolium</i> (Tourn.) L.	
* <i>Trifolium arvense</i> L.	Rabbit-foot or Stone
Dry sandy soil, fields, road-	Clover
sides. North Takoma 1912.	May-July
<i>Native of Europe.</i>	
<i>Trifolium pratense</i> L.	Red Clover
Abundant in fields and meadows.	May-August
Introduced from Europe. C.	
<i>Trifolium repens</i> L.	White Clover
Fields, etc.; abundant.	May-August
<i>Trifolium hybridum</i> L.	Alsike Clover
Fields, etc.; frequent.	May-September
European.	
* <i>Trifolium agrarium</i> L.	Yellow or Hop Clover
<i>Meibomia</i> Fields, etc.; frequent.	June-August
Beyond Soldiers Home 1896.	
European.	
<i>Melilotus</i> (Tourn.) Hill	Melilot. Sweet Clover
<i>Melilotus officinalis</i> (L.) Lam.	Yellow Melilotus
Frequent in waste ground, etc.	May-August
Native of Europe and Asia.	
<i>Melilotus alba</i> Dear.	White Sweet Clover
Waste ground, etc.; frequent.	May-August
Native of Europe.	
<i>Tephrosia</i> Pers.	Hoary Pea
<i>Cracca</i> L. (L.) Britton	
Dry woods; occasional.	
* <i>Tephrosia virginiana</i> (L.) Pers.	Goat's Rue, Catgut
<i>Cracca virginiana</i> L. (L.) Britton	
Takoma 1895.	May-June
<i>Robinia</i> L. Hornem.	Locust
Dry woods and woods; infrequent.	
<i>Robinia pseudo-acacia</i> L. Common L.	False Acacia. Black
Abundant.	Locust.
<i>Desmodium</i> Desv.	May
<i>Meibomia</i> Heist.	
<i>Desmodium nudiflorum</i> (L.) D C.	Tick Trefoil
<i>Meibomia nudiflora</i> (L.) Kuntze	
Deciduous woods; common.	July-September

* <i>Desmodium grandiflorum</i> (Walt.) D C.	Purple Flower
<i>Meibomia grandiflora</i> (Walt.) Kuntze	
Occasional in woods. B.S.P.	June-July
Kensington 1910.	July-September
* <i>Desmodium viridiflorum</i> (L.) Beck	
<i>Meibomia viridiflora</i> (L.) Kuntze	
Infrequent in woods. Slighter but	June-August
Terra Cotta 1894.	August-September
<i>Desmodium Dillenii</i> Darl.	Vetch, Terra.
<i>Meibomia dillenii</i> (Darl) Kuntze	
Occasional in woods. Orchard.	August-September
<i>Desmodium paniculatum</i> (L.) D C.	April-June
<i>Meibomia paniculata</i> (L.) Kuntze	
Occasional in woods and thickets.	August-September
* <i>Desmodium obtusum</i> (Muhl.) D C.	May-July
<i>Meibomia obtusa</i> (Muhl.) Vail.	
Occasional in dry woods, hills, etc. Takoma 1910.	August-September
<i>Viola pedunculata</i> (L.) Moench.	Hairy or Stalked
* <i>Desmodium marilandicum</i> (L.) D C.	May-September
<i>Meibomia marilandica</i> (L.) Kuntze	
Frequent in dry woods, etc.	August-September
<i>Lespedeza</i> Michx.	Bush Clover
<i>Lespedeza procumbens</i> Michx.	Ground Cover
Dry fields and woods; frequent.	August-September
<i>Apis tuberosa</i> (L.) Moench.	
<i>Lespedeza repens</i> (L.) Bartr.	
Dry fields and woods; common.	July-September
* <i>Lespedeza Nuttallii</i> Darl.	Kidney Bean
Dry woods; not frequent.	August-September
Takoma 1905.	Wild Bean
<i>Lespedeza virginica</i> (L.) Britton	August-September
Dry woods; occasional.	August-October
* <i>Lespedeza frutescens</i> (L.) Britton	
Occasional. Takoma 1905.	August-September
* <i>Lespedeza hirta</i> (L.) Hornem.	Butterfly Plant
Dry fields and woods; infrequent.	August-October
Woodside 1896.	
<i>Lespedeza striata</i> (Thunb.) Hook and Arn.	June-August
Dry fields and woods; frequent.	Japan Clover
Native of East Asia.	August-October

	<i>Stylosanthes</i> Swartz	Pencil Flower.
	<i>Stylosanthes biflora</i> (L.) B.S.P. Dry soil; frequent.	July-September
	* <i>Stylosanthes riparia</i> Kearney Dry soil; occasional. Takoma 1895, labeled <i>elatior</i> but undoubtedly <i>riparia</i> .	August-September June-August
	<i>Vicia</i> (Tourn.) L. Dry soil.	Vetch. Tare. July-September
	* <i>Vicia angustifolia</i> (L.) Reichard. Fields and waste ground; occasional. Native of Europe. Silver Spring 1915.	Common Vetch April-June
	* <i>Vicia tetrasperma</i> (L.) Moench. Fields, etc.; infrequent. Native of Europe. Below Takoma 1916.	Common Vetch May-July
	<i>Vicia villosa</i> Roth. Sometimes cultivated and common, escaping. Native of Europe and Asia. Fields, etc.; occasionally. Witskie 1938.	Hairy or Winter Vetch May-September
	<i>Apios</i> (Boerh.) Ludwig. <i>Glycine</i> L.	Groundnut. Wild Bean June-August
OXALIDACEAE (WATER-PEA FAMILY)		
	<i>Apios tuberosa</i> (L.) Moench. <i>Glycine apios</i> L. Wet places; infrequent.	Food Berry July-September
	<i>Phaseolus</i> (Tourn.) L.	Kidney Bean
	<i>Phaseolus polystachyus</i> (L.) B.S.P. Waste ground, low thickets; rare.	Wild Bean August-September
	<i>Strophostyles</i> Ell.	Wild Bean
	<i>Strophostyles umbellata</i> (Muhl.) Britton Occasional.	July-August
	<i>Clitoria</i> L.	Butterfly Pea
	* <i>Clitoria mariana</i> L. Dry or moist ground; occasional. Terra Cotta 1895.	Wild Gram June-August
	dry fields, rocky poor soil. Terra Cotta 1895.	April-May

POLYGALACEAE (Amphicarpa Ell. Falcata Gmel.)	Hog Peanut Milk-vetch
*Amphicarpa monoica (L.) Ell. Falcata comosa (L.) Kuntze Moist thickets; occasional. Kensington 1910, Takoma 1912.	August-September
*Polygala Galactia P. Br. Polygala volubilis L. Galactis volubilis (L.) Britton Dry soil; frequent.	Milk Pea July-September July-September

LINACEAE (FLAX FAMILY)	July-September
Dry sandy soil. Takoma 1903. Linum (Tourn.) L.	Flax
*Polygala Contorta Gray. *Linum usitatissimum L. Cultivated flax infrequently escaped. European.	Common Flax
*Polygala contorta Gray. Woodside 1896. Infrequent. Rock Creek 1897. Rock Creek	July-September
*Linum striatum Walt. Swamps, wet places; not common.	July-September
*Polygala Brightwood 1896 in Rock Creek valley. Dry fields, etc.; occasional.	July-September
*Linum medium (Planch.) Britton Dry soil; infrequent.	June-August
Takoma 1897.	

EUPHORBIACEAE (JADE PLANT)	Three-angled Jade
*Euphorbia Oxalis L. Xanthoxalis Small.	Wood Sorrel Yellow Wood Sorrel

*Oxalis filipes Small. Xanthoxalis filipes Small. Woods; common. Takoma 1904.	Spurge May-August
Oxalis corniculata L. Xanthoxalis corniculata (L.) Small. Rare. Dry fields, where grown.	Lady's Sorrel June-September

GERANIACEAE (GERANIUM FAMILY)	Wild Cranesbill
Geranium maculatum L. Geranium (Tourn.) L.	Cranesbill
Geranium maculatum L. Woods; frequent.	Wild Cranesbill April-June
*Geranium carolinianum L. Dry fields, rocky poor soil.	July-October
Terra Cotta 1895.	April-May
Geranium maculatum L. The only gerani.	Water Geranium

POLYGALACEAE (MILKWORT FAMILY)

In shallow water in marsh near
mouth Polygala (Tourn.) L.

Milkwort

**Polygala incarnata* L.

Dry sandy soil; infrequent.
Woodside 1895.

July-September

**Polygala sanguinea* L.

Polygala viridescens L.

Sandy and moist ground.
Woodside 1895.

June-October

**Polygala Nuttallii* T. and G.

Dry sandy soil. Takoma 1903.

July-September

**Polygala Curtissii* Gray.

Takoma 1910.

July-October

**Polygala cruciata* L.

Swamps, etc.; infrequent.
Brightwood 1897. Rock Creek to
drainage. By touch or contact
will dry and then burn.

July-September

**Polygala verticillata* L.

Dry soil, fields, etc.; occasional.
Woodside 1896.

July-October

EUPHORBIACEAE (SPURGE FAMILY)

To many species very poisonous to
humans.

Three-seeded Mercury

**Acalypha virginica* L.

Fields, gardens, etc.; occasional. June-October
Terra Cotta 1895.

Spurge

Euphorbia L.

Chamaesyce S. F. Gray

Euphorbia Preslii Guss

Chamaesyce preslii (Guss.) Arth.

Cultivated fields, waste ground.

May-October

Euphorbia maculata L.

Milk purslane

Chamaesyce maculata (L.) Small

Dry open ground, walks; common.

June-November

Euphorbia corollata L.

Flowering Spurge

Tithymalopsis corollata (L.) Klotzsch and Gareke

Dry open places; frequent.

July-October

CALLITRICHACEAE (WATER STARWORT FAMILY)

Callitricha L.

Water Starwort

The only genus.

CEL **Callitricha palustris* L.
In shallow water in marsh near
mouth of Sligo.
Takoma 1904.

**Callitricha heterophylla* Pursh.
In shallow water; Woodside 1895.

Fr. May-August

Fr. May-August

ANACARDIACEAE (CASHEW FAMILY)

Rhus L.

Sumach

Rhus glabra L. March.
Dry soil; common. Planted

Smooth or White Sumac
July

*Rhus copallina L.
Dry places; frequent.
Woodside 1895.

Dwarf Sumac
July

Rhus vernix L.
Toxicodendron vernix (L.) Kuntze
Swamps, etc. Very poisonous to
many people by touch or contact
with the smoke when burned.
Common.

Poison Dogwood
Poison Sumac
Poison Elder
June-July

Rhus Toxicodendron L.
Toxicodendron radicans (L.) Kuntze
Abundant everywhere. To many
persons violently poisonous to
the touch.

Poison Ivy. Poison
Oak.
May-June

AQUIFOLIACEAE (HOLLY FAMILY)

Cleopatra Ilex L.

Holly

Ilex opaca Ait.
Low woods; infrequent.

American Holly
May-June; fr. August-
September

Ilex decidua Walt.
Moist woods; rare.
Terra Cotta 1876.

Deciduous Holly
May-June; fr. August-
September

Ilex verticillata (L.) Gray
Low ground; rare.
Terra Cotta 1878.

Black Alder. Winter-
berry.
June; fr. August

Ilex laevigata (Pursh.) Gray
Wet ground; very rare.
Terra Cotta 1882.

Smooth Winterberry
May-June; fr. August-
September

CELASTRACEAE (STAFF TREE FAMILY. BITTERSWEET FAMILY)

Euonymus (Tourn.) L. Spindle Tree

**Euonymus americanus* L. Strawberry Bush
Wooded river bank; occasional. May
Sligo below Takoma 1915.

ACERACEAE (MAPLE FAMILY)

Acer sp. spicatum L. Acer (Tourn.) L. Maple
Waste places

Acer saccharum Marsh. Rock or Sugar Maple
Rich woods; rare. Planted May
along streets, etc.

Acer saccharinum L. White or Silver Maple
River banks; frequent. January-March

Acer rubrum L. Red or Swamp Maple
Abundant February-April

BALSAMINACEAE (TOUCH-ME-NOT FAMILY)

Impatiens (Rivinius) L. Balsam. Jewelweed

Impatiens biflora Walt. Spotted Touch-me-not
Rills and wet shady places; June-September
common. soil; infrequent.

RHAMNACEAE (BUCKTHORN FAMILY)

Clethra sp. spicata L. Red-root

Clethra alnifolia L. New Jersey Tea
Open woods. Tea substitute
during the Revolutionary War.

VITACEAE (VINE FAMILY)

Pedera Neck. Virginia Creeper.
Parthenocissus Planch. Woodbine

Parthenocissus quinquefolia (L.) Planch. Virginia Creeper
(*Pedera quinquefolia* (L.) Greene)
Deciduous woods; common. May-June; fr. August-September

Vitis (Tourn.) L. Grape
Thickets, etc. River Bank or Frost
Terra Cotta 1895. Grape.
May-June; fr. July-August,

MALVACEAE (MALLOW FAMILY)

<i>Abutilon</i> (Tourn.) Mill.	Indien Mallow
<i>Abutilon Theophrasti</i> Medic. Wastes, fields; frequent	Velvet Leaf July-August
<i>Sida L.</i>	Firwood
<i>Sida spinosa</i> L. Wastes, etc.	False Mallow July-September
<i>Malva</i> (Tourn.) L.	Mallow
* <i>Malva rotundifolia</i> L. Wastes; frequent. Takoma 1896. Native of Europe and Asia.	Common M., Cheeses. May-August
<i>Hibiscus L.</i>	Rose Mallow
<i>Hibiscus syriacus</i> L. Cultivated and escaping.	Shrubby Althea. Rose of Sharon.

HYPERICACEAE (ST. JOHN'S-WORT FAMILY)

<i>Ascyrum L.</i>	St. Peter's-wort
* <i>Ascyrum stans</i> Michx. Dry sandy soil; infrequent. Brightwood 1897.	St. Peter's-wort August
* <i>Ascyrum hypericoides</i> L. Sandy rocky places; occasional. Woodside 1895.	St. Andrew's Cross July-September
<i>Hypericum (Tourn.) L.</i>	St. John's-wort
* <i>Hypericum perforatum</i> L. Fields, etc.; occasional. Woodside 1896.	Common St. John's-wort July
* <i>Hypericum punctatum</i> Lam. <i>Hypericum denticulatum</i> Walt. Wet places. Sligo two miles below Takoma 1895. Labelled <i>Maculatum</i> L.	July
<i>Hypericum mutilum</i> L. Wet places; occasional.	August-October
* <i>Hypericum gentianoides</i> (L.) B.S.P. <i>Sarcandra gentianoides</i> Sandy or rocky soil. Woodside 1895.	Orange Grass. Pine- weed. August-September